

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# **Product datasheet for TA503830**

## **NUDT18 Mouse Monoclonal Antibody [Clone ID: OTI1A5]**

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI1A5

**Applications:** FC, IF, IHC, WB

**Recommend Dilution:** WB 1:2000, IHC 1:150, IF 1:100, FLOW 1:100

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human NUDT18(NP\_079091) produced in

HEK293T cell.

**Formulation:** PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

**Concentration:** 0.6 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Predicted Protein Size: 35.3 kDa


**Gene Name:** nudix hydrolase 18

Database Link: NP 079091 Entrez Gene 79873 Human


Synonyms: MTH3



## **Product images:**




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY NUDT18 ([RC204376], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-NUDT18. Positive lysates [LY403033] (100ug) and [LC403033] (20ug) can be purchased separately from OriGene.




Immunohistochemical staining of paraffinembedded Human breast tissue within the normal limits using anti-NUDT18 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA503830)






Immunohistochemical staining of paraffinembedded Human pancreas tissue within the normal limits using anti-NUDT18 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA503830)




Immunohistochemical staining of paraffinembedded Human endometrium tissue within the normal limits using anti-NUDT18 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA503830)




Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human endometrium tissue using anti-NUDT18 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA503830)





Anti-NUDT18 mouse monoclonal antibody (TA503830) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY NUDT18 ([RC204376]).



HEK293T cells transfected with either [RC204376] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-NUDT18 antibody (TA503830), and then analyzed by flow cytometry.